

Analytical Solution of the $SL(2, \mathbb{R})/U(1)$ WZNW Black Hole Model

Uwe Müller* and Gerhard Weigt**

Deutsches Elektronensynchrotron DESY
Institut für Hochenergiephysik IfH Zeuthen
Platanenallee 6, D-15738 Zeuthen, Germany

Abstract

The gauged $SL(2, \mathbb{R})/U(1)$ Wess-Zumino-Novikov-Witten (WZNW) model is classically an integrable conformal field theory. We have found a Lax pair representation for the non-linear equations of motion, and a Bäcklund transformation. A second-order differential equation of the Gelfand-Dikii type defines the Poisson bracket structure of the theory, and its fundamental solutions describe the general solution of the WZNW model as well. The physical and free fields are related by non-local transformations. The (anti-)chiral component of the energy-momentum tensor which factorizes into conserved quantities satisfying a non-linear algebra characteristic for parafermions assumes the canonical free field form. So the black hole model is expected to be prepared for an exact canonical quantization.

Witten has shown [1] that the conformal $SL(2, \mathbb{R})/U(1)$ gauged WZNW action [2] describes string propagation in the background of a two-dimensional space-time black hole. He conjectured that “the conformal field theory governing the black hole is more or less exactly soluble.” However, this theory is not conformal in the ordinary sense. It is true that it has classically a traceless energy-momentum tensor, but the β -function is non-zero [3, 1, 4, 5] as the gauged WZNW action corresponds to a σ -model with curvature. The inherent non-local structure of the theory might settle this question. But that requires, obviously, a complete analytical solution of the theory, and

*E-mail: umueller@ifh.de

**E-mail: weigt@ifh.de

its exact quantization. Classical solutions were found for two-dimensional string theory in any curved space-time [6]. But we are looking here for unconstrained field-theoretic solutions. Our intention is to find a canonical transformation of the physical fields onto free fields as a prerequisite for a canonical quantization of the present non-linear theory.

Interestingly, there is an embedding of the black hole in an integrable non-abelian Toda theory [7]. Although this apparently does not solve the addressed problem the results of refs. [7] and [8] stimulated the analytical calculations presented in this letter. We could show that the analytical solution of the $SL(2, \mathbb{R})/U(1)$ coset theory is asymptotically related to the solution of that non-abelian Toda theory [9]. However, there is an independent way of proving integrability of the $SL(2, \mathbb{R})/U(1)$ model. We have found a Lax pair representation for the non-linear equations of motion, and a Bäcklund transformation as well. We have solved the equations in general, and derived a non-local transformation of the interacting fields onto free fields for periodic boundary conditions.

In this letter we discuss the classical theory. Here we will not be able to give all the details and proofs, in particular, we shall omit the Lax pair representation completely and refer to a separate paper [10].

The $SL(2, \mathbb{R})/U(1)$ gauged WZNW action can be derived from the $U(1)$ gauge invariant WZNW action. In light-cone coordinates $z = \tau + \sigma$, $\bar{z} = \tau - \sigma$, $\partial_z = (\partial_\tau + \partial_\sigma)/2$, $\partial_{\bar{z}} = (\partial_\tau - \partial_\sigma)/2$ this action is ($\gamma \equiv \sqrt{2\pi/k}$)

$$\begin{aligned}
S[g, A] = & \frac{1}{\gamma^2} \int_M \left[\frac{1}{2} \text{tr} (g^{-1} \partial_z g g^{-1} \partial_{\bar{z}} g) + \text{tr} (I g^{-1} \partial_z g) A_{\bar{z}} + \right. \\
& \left. + \text{tr} (I \partial_{\bar{z}} g g^{-1}) A_z + (\text{tr} (I g I g^{-1}) - 2) A_z A_{\bar{z}} \right] dz d\bar{z} + \\
& + \frac{1}{6\gamma^2} \int_{B, \partial B=M} \text{tr} (g^{-1} dg \wedge g^{-1} dg \wedge g^{-1} dg). \tag{1}
\end{aligned}$$

Here I is a matrix

$$I = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \tag{2}$$

and the Lie group-valued field g is parameterized by (σ_3 is the third Pauli matrix)

$$g = g(r, t, \alpha) = \exp((\alpha + t)I/2) \exp(r\sigma_3) \exp((\alpha - t)I/2). \tag{3}$$

In order to define the gauged WZNW action we have to eliminate the abelian gauge field $A_z, A_{\bar{z}}$. This will not be done by path integration [11, 1, 5].

Such integrations are still incomplete because functional determinants remain uncalculated [11], and we would not get a suitable effective action to start with. Instead, we eliminate the gauge field through its equation of motion and gauge the angle variable α by $\alpha = 0$. So we obtain an exact classical action [2]

$$S[r, t] = \frac{1}{\gamma^2} \int_M (\partial_z r \partial_{\bar{z}} r + \tanh^2 r \partial_z t \partial_{\bar{z}} t) dz d\bar{z}. \quad (4)$$

Its (euclidean signature) black hole [1] target-space metric

$$ds^2 = dr^2 + \tanh^2 r dt^2 \quad (5)$$

has the form of a semi-infinite cigar which is asymptotic for $r \rightarrow \infty$ to $\mathbb{R} \times S^1$ with a flat metric. The conformal invariance of this theory is indicated by the tracelessness of its energy-momentum tensor.

The equations of motion following from (4) are

$$\begin{aligned} \partial_z \partial_{\bar{z}} r &= \frac{\sinh r}{\cosh^3 r} \partial_z t \partial_{\bar{z}} t, \\ \partial_z \partial_{\bar{z}} t &= -\frac{1}{\sinh r \cosh r} (\partial_z r \partial_{\bar{z}} t + \partial_z t \partial_{\bar{z}} r). \end{aligned} \quad (6)$$

They provide conservation of the chiral component of the energy-momentum tensor (we will not indicate, whenever possible, the similar anti-chiral parts)

$$T = \frac{1}{2} (T_{\tau\tau} + T_{\tau\sigma}) = \frac{1}{\gamma^2} ((\partial_z r)^2 + \tanh^2 r (\partial_z t)^2). \quad (7)$$

Moreover, there are two further conserved chiral quantities on shell

$$V_{\pm} = \frac{1}{\gamma^2} e^{\pm i\nu} (\partial_z r \pm i \tanh r \partial_z t). \quad (8)$$

For the conservation law $\partial_{\bar{z}} V_{\pm} = 0$, the (non-local) ν is sufficiently defined by the derivatives [7]

$$\partial_z \nu = (1 + \tanh^2 r) \partial_z t, \quad \partial_{\bar{z}} \nu = \cosh^{-2} r \partial_{\bar{z}} t. \quad (9)$$

However, it will be easy to integrate them, once we have got the general solution of the equations of motion, since the integrability conditions of (9) just yield the second equation of (6). Surprisingly, the chiral component of the energy-momentum tensor factorizes in terms of the $V_{\pm}(z)$ and looks like a Sugawara construction

$$T = \gamma^2 V_+ V_-. \quad (10)$$

But the conformal spin-one quantities $V_{\pm}$ are no usual Kac-Moody currents.

The general solution of the equations of motion will be stated here without proof [10]. It can easily be seen that

$$\begin{aligned}\sinh^2 r &= X\bar{X}, \\ t &= i(B - \bar{B}) + \frac{i}{2} \ln \frac{X}{\bar{X}}\end{aligned}\tag{11}$$

solve the equations of motion (6) if

$$\begin{aligned}X &= A + \frac{\bar{B}'}{\bar{A}'}(1 + A\bar{A}), \\ \bar{X} &= \bar{A} + \frac{B'}{A'}(1 + A\bar{A}).\end{aligned}\tag{12}$$

$A(z), B(z), A'(z), B'(z)$ ($\bar{A}(\bar{z}), \bar{B}(\bar{z}), \bar{A}'(\bar{z}), \bar{B}'(\bar{z})$) are arbitrary chiral (anti-chiral) functions and their derivatives, respectively. The solution (11, 12) is invariant under $GL(2, \mathbb{C})$ transformations

$$\begin{aligned}A &\rightarrow T[A] = \frac{aA - b}{cA + d}, \\ B &\rightarrow T[B] = B + \ln(cA + d), \\ \bar{A} &\rightarrow T[\bar{A}] = \frac{d\bar{A} - c}{b\bar{A} + a}, \\ \bar{B} &\rightarrow T[\bar{B}] = \bar{B} + \ln(b\bar{A} + a), \\ &\left(\begin{array}{cc} a & -b \\ c & d \end{array} \right) \in GL(2, \mathbb{C}).\end{aligned}\tag{13}$$

In order to find free field realizations of these functions, first we have to calculate their Poisson brackets using the canonical Poisson brackets of the physical fields r, t (at equal τ !). The procedure applied in ref. [8] is an indirect one and does not prove a unique result. We shall calculate the Poisson brackets directly by using the characteristic differential equation

$$y'' - (\partial_z V_- / V_-)y' - \gamma^2 T y = 0,\tag{14}$$

which is derived from the conserved quantities (8). (We should mention here that this equation fits into the Gelfand-Dikii hierarchy¹ with $W^{(1)} = (\partial_z V_- / V_-)$ and that it could be transformed with a modified energy-momentum tensor $\tilde{T}$ into

$$\tilde{y}'' - \tilde{T}\tilde{y} = 0,\tag{15}$$

¹We thank Adel Bilal for discussions about the Gelfand-Dikii hierarchy.

well-known in Liouville theory. y has conformal weight zero, $\tilde{y} = y/\sqrt{V_-}$ conformal weight $-1/2$.) To get $A(z)$, $B(z)$ related to the solutions of this equation, we have to integrate the eqs. (9) so that the $V_{\pm}$ become defined. From the general solution (11, 12) one obtains

$$\nu - t = i(B + \bar{B}) - \frac{i}{2} \ln \frac{1 + X\bar{X}}{(1 + A\bar{A})^2}, \quad \nu + \bar{\nu} = 2t. \quad (16)$$

We have, furthermore, to rewrite the coefficients of eq. (14) completely in terms of $A(z)$, $B(z)$ using again (11, 12). The two fundamental solutions of eq. (14) are then given by

$$\begin{aligned} y_1(z) &= e^{B(z)}, \\ y_2(z) &= y_1(z) \int^z \frac{W(z')}{y_1^2(z')} dz'. \end{aligned} \quad (17)$$

The Wronski determinant

$$W = y_1 y_2' - y_2 y_1' \quad (18)$$

is here related by means of eq. (14) to V_-

$$W = \gamma^2 V_- = A' e^{2B}, \quad (19)$$

so that we find $A(z)$ and $B(z)$ explicitly in terms of the fundamental solutions

$$A(z) = \frac{y_2}{y_1}, \quad B(z) = \ln y_1. \quad (20)$$

The Poisson brackets of the $A(z)$, $B(z)$ can be calculated once we have got those of the y_1 , y_2 , and, finally, we could read off the $A(z)$ and $B(z)$ as functions of free fields.

The differential equation (14) determines the dependence of the y_i on the fields r , t and their momenta π_r , π_t . In order to derive the Poisson brackets, we need the variations δy_i in terms of the variations δr , δt , $\delta \pi_r$, and $\delta \pi_t$, which enter the definition of the Poisson bracket. Varying eq. (14) we obtain

$$\delta y'' - (\partial V_- / V_-) \delta y' - \gamma^2 T \delta y = \delta (\partial V_- / V_-) y' + \gamma^2 \delta T y. \quad (21)$$

The homogeneous part of this differential equation is identical to eq. (14). To find a unique solution of the inhomogeneous equation we need subsidiary conditions. Using field-theoretic initial conditions (no zero modes!)

$\delta y_i(-\infty) = 0$, $\delta y'_i(-\infty) = 0$, it can be represented by the two solutions y_1 and y_2 of the homogeneous equation

$$\begin{aligned}\delta y_i(\sigma) &= \int_{-\infty}^{\sigma} d\sigma' \Omega(\sigma, \sigma') \left(\delta(\partial V_-/V_-)(\sigma') y'_i(\sigma') + \gamma^2 \delta T(\sigma') y_i(\sigma') \right), \\ \Omega(\sigma, \sigma') &\equiv \frac{y_1(\sigma') y_2(\sigma) - y_2(\sigma') y_1(\sigma)}{y_1(\sigma') y'_2(\sigma') - y_2(\sigma') y'_1(\sigma')}.\end{aligned}\quad (22)$$

This leads directly to the Poisson brackets of the y_i [10]

$$\{y_i(\sigma), y_j(\tilde{\sigma})\} = \frac{\gamma^2}{2} (y_i(\sigma) y_j(\tilde{\sigma}) - y_j(\sigma) y_i(\tilde{\sigma})) \epsilon(\sigma - \tilde{\sigma}). \quad (23)$$

$\epsilon(\sigma)$ denotes here the sign function.

The Poisson brackets for the $A(z)$, $B(z)$ are found by taking into consideration eqs. (20), and we read off the free field realizations [10]

$$\begin{aligned}A'(z) &= \gamma (\partial_z \phi_1 - i \partial_z \phi_2) \exp(-2\gamma \phi_1(z)), \\ B(z) &= \gamma (\phi_1 - i \phi_2).\end{aligned}\quad (24)$$

The conserved $V_{\pm}(z)$ then become

$$V_{\pm}(z) = \frac{1}{\gamma} (\partial_z \phi_1 \pm i \partial_z \phi_2) \exp(\pm 2i \gamma \phi_2(z)), \quad (25)$$

and, as expected, the energy-momentum tensor T assumes the canonical free field form

$$T = (\partial_z \phi_1)^2 + (\partial_z \phi_2)^2. \quad (26)$$

Vice versa, we could get the same free field realizations (24), consistently, as solutions of eq.(14) if its coefficients would have been given by (25, 26), and, third, by requiring that the free fields of the energy-momentum tensor (26) are local in $y_i(z)$ and their derivatives [10]. This remains true for the periodic string case, after including the corresponding zero modes. But for periodic boundary conditions the non-periodic sign function has to be replaced by the periodic sawtooth function

$$h(\sigma) = \epsilon(\sigma) - \frac{1}{\pi} \sigma, \quad (27)$$

and we have to take into consideration the periodic δ -function

$$\delta_{2\pi}(\sigma) \equiv \frac{1}{2} \partial_{\sigma} \epsilon(\sigma) = \sum_n \delta(\sigma - 2\pi n). \quad (28)$$

$\epsilon(\sigma)$ denotes now the stair-step function

$$\epsilon(\sigma) = 2n + 1 \quad \text{for} \quad 2n\pi < \sigma < (2n + 1)\pi, \quad (29)$$

which coincides with $\text{sign}(\sigma)$ for $-2\pi < \sigma < 2\pi$. (The principal value is understood at the discontinuities.)

It is worth mentioning here that the conserved $V_{\pm}(z)$ are related to the $\text{SL}(2, \mathbb{R})/\text{U}(1)$ coset currents. They satisfy non-linear Poisson brackets

$$\begin{aligned} \{V_{\pm}(\sigma), V_{\pm}(\sigma')\} &= \gamma^2 V_{\pm}(\sigma) V_{\pm}(\sigma') \epsilon(\sigma - \sigma'), \\ \{V_{\pm}(\sigma), V_{\mp}(\sigma')\} &= -\gamma^2 V_{\pm}(\sigma) V_{\mp}(\sigma') \epsilon(\sigma - \sigma') + \frac{1}{\gamma^2} \delta'(\sigma - \sigma') \end{aligned} \quad (30)$$

which are characteristic for parafermions [12, 2].² This non-linear algebra also provides the Virasoro algebra, and implies conformal weight one for the $V_{\pm}$

$$\{T(\sigma), V_{\pm}(\sigma')\} = -(\partial_{\sigma'} V_{\pm}(\sigma') \delta(\sigma - \sigma') - V_{\pm}(\sigma') \delta'(\sigma - \sigma')). \quad (31)$$

It remains to calculate $A(z)$ by integrating $A'(z)$. We choose now (closed string) periodic boundary conditions, and in the target space

$$\begin{aligned} r(\tau, \sigma + 2\pi) &= r(\tau, \sigma), \\ t(\tau, \sigma + 2\pi) &= t(\tau, \sigma) + 2\pi w, \quad w \in \mathbb{Z}. \end{aligned} \quad (32)$$

As $t(z)$ is an angle variable in the parameterization (3), w is a ‘winding number’ describing how often the string surrounds the origin of the target space. In the following we restrict our general solution to real free fields $\psi_i(\tau, \sigma) = \phi_i(z) + \bar{\phi}_i(\bar{z})$ and find in accordance with (32) the periodicity properties

$$\psi_i(\tau, \sigma + 2\pi) = \psi_i(\tau, \sigma) + \frac{2\pi w}{\gamma} \delta_{i,2}, \quad (33)$$

which assign ψ_1 and ψ_2 directly to $r(z)$ respectively $t(z)$. The corresponding mode expansions are given by

$$\begin{aligned} \phi_i(z) &= \frac{1}{2} q_i + \left(\frac{1}{4\pi} p_i + \frac{w}{2\gamma} \delta_{i,2} \right) z + \frac{i}{\sqrt{4\pi}} \sum_{n \neq 0} \frac{a_n^{(i)}}{n} e^{-inz}, \\ \bar{\phi}_i(\bar{z}) &= \frac{1}{2} q_i + \left(\frac{1}{4\pi} p_i - \frac{w}{2\gamma} \delta_{i,2} \right) \bar{z} + \frac{i}{\sqrt{4\pi}} \sum_{n \neq 0} \frac{\bar{a}_n^{(i)}}{n} e^{-in\bar{z}}. \end{aligned} \quad (34)$$

²We thank Konstadinos Sfetsos for a comment on parafermions in this respect. Instead of solving the present conformal theory by an infinite set of conservation laws, we are intending to emphasize the dynamics of the theory governed by the general solutions of its equations of motion.

This tells us that $A(z)$ is periodic up to a factor (comp. (24))

$$A(z + 2\pi) = \exp(-\gamma p_1) A(z). \quad (35)$$

The integration of $A'(z)$ then yields $A(z)$ non-locally related to the free fields

$$A(z) = -\frac{\gamma}{2} \sinh^{-1}\left(\frac{\gamma p_1}{2}\right) \times \int_0^{2\pi} dz' \exp\left(-\frac{\gamma p_1}{2} \epsilon(z - z')\right) (\partial_{z'} \phi_1 - i \partial_{z'} \phi_2) \exp(-2\gamma \phi_1(z')). \quad (36)$$

With

$$\partial_z \exp(\alpha \epsilon(z - z')) = 2 \sinh \alpha \delta_{2\pi}(z - z') \exp\left(\frac{\alpha}{\pi} (z - z')\right), \quad (37)$$

we get back the local $A'(z)$ of (24). As seen from (16) the conserved $V_{\pm}(z)$ are not periodic, except the zero mode of the angle variable ψ_2 given by $\gamma p_2 = \int_0^{2\pi} dz' \dot{t} \tanh^2 r$ becomes discrete $\gamma p_2 = 2\pi n$, $n \in \mathbb{Z}$.

The general solution of the dynamical eqs. (6) is only complete after we have included the corresponding anti-chiral parts. Choosing different target-space coordinates

$$u = \sinh r e^{it}, \quad \bar{u} = \sinh r e^{-it}, \quad (38)$$

the (covariantly written) action (4) becomes

$$S = \frac{1}{2\gamma^2} \int d\tau d\sigma \sqrt{-g} g^{ij} \frac{\partial_i u \partial_j \bar{u}}{1 + u \bar{u}}. \quad (39)$$

It shows for Wick rotated t the metric singularity of a black hole [1]. In terms of these new coordinates the solution (11, 12) gets a rather simple form if it is expressed by $y_1(z)$, $y_2(z)$ (20) (respectively $\bar{y}_1(\bar{z})$, $\bar{y}_2(\bar{z})$)

$$u = \frac{\bar{y}_1 y_1' + \bar{y}_2 y_2'}{y_1 y_2' - y_1' y_2}, \quad \bar{u} = \frac{y_1 \bar{y}_1' + y_2 \bar{y}_2'}{\bar{y}_1 \bar{y}_2' - \bar{y}_1' \bar{y}_2}. \quad (40)$$

The $\text{GL}(2, \mathbb{C})$ invariance of the solutions now translates into

$$\begin{pmatrix} y_1 \\ y_2 \end{pmatrix} \rightarrow \begin{pmatrix} d & c \\ -b & a \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \quad \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \end{pmatrix} \rightarrow \begin{pmatrix} a & b \\ -c & d \end{pmatrix} \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \end{pmatrix}. \quad (41)$$

Introducing the complex free fields

$$\psi = \psi_1 + i\psi_2, \quad \bar{\psi} = \psi_1 - i\psi_2, \quad (42)$$

we find a Bäcklund transformation of the equations of motion of the $u, \bar{u}$

$$\begin{aligned}\partial_z u &= \frac{\gamma}{2} e^{\gamma\psi} \left(P + i\sqrt{4Q - P^2} \right) \partial_z \psi, & \partial_{\bar{z}} u &= \frac{\gamma}{2} e^{\gamma\psi} \left(P - i\sqrt{4Q - P^2} \right) \partial_{\bar{z}} \psi, \\ \partial_z \bar{u} &= \frac{\gamma}{2} e^{\gamma\bar{\psi}} \left(P - i\sqrt{4Q - P^2} \right) \partial_z \bar{\psi}, & \partial_{\bar{z}} \bar{u} &= \frac{\gamma}{2} e^{\gamma\bar{\psi}} \left(P + i\sqrt{4Q - P^2} \right) \partial_{\bar{z}} \bar{\psi},\end{aligned}\tag{43}$$

where

$$P(u, \bar{u}, \psi, \bar{\psi}) = u e^{-\gamma\psi} + \bar{u} e^{-\gamma\bar{\psi}} + e^{-\gamma\psi - \gamma\bar{\psi}}, \quad Q(u, \bar{u}, \psi, \bar{\psi}) = (1 + u\bar{u}) e^{-\gamma\psi - \gamma\bar{\psi}}.\tag{44}$$

The integrability conditions for the fields u and $\bar{u}$ are equivalent to the free field equations for ψ and $\bar{\psi}$, and those for the ψ and $\bar{\psi}$ provide the non-linear equations of motion. The Bäcklund transformation (43, 44) respects periodicity.

Although the classical solution of the $SL(2, \mathbb{R})/U(1)$ gauged WZNW model (4) bears strong resemblance to Liouville or Toda theories its quantum structure might be different because its energy-momentum tensor does not have a ‘central charge’ term classically. Furthermore, we would like to see whether the black hole singularity survives quantization, a dilaton might appear, and whether the Hawking radiation could be discussed in some manner based on this analytical solution.

References

- [1] E. Witten, *Phys. Rev.* **D44** (1991) 314.
- [2] K. Bardakci, M. Crescimanno, E. Rabinovici, *Nucl. Phys.* **344** (1990) 344.
- [3] C.G. Callan, D. Friedan, E.J. Martinec, and M.J. Perry, *Nucl. Phys.* **B262** (1985) 593.
- [4] R. Dijkgraaf, E. Verlinde, H. Verlinde, *Nucl. Phys.* **B371** (1992) 269.
- [5] A. A. Tseytlin, *Nucl. Phys.* **B399** (1993) 601.
- [6] I. Bars, J. Schulze, *Phys. Rev.* **D51** (1995) 1854; I. Bars, Lectures given at 2nd Journee Cosmologique, Paris, France, 2-4 June 1994, published in Journee Cosmologique 1994, 399-428, hep-th/9411217.
- [7] J.-L. Gervais, M. Saveliev, *Phys. Lett.* **B286** (1992) 271.

- [8] A. Bilal, *Nucl. Phys.* **B422** (1994) 258.
- [9] To appear in the Proceedings of the 30th International Symposium Ahrenshoop, August 26 – 31, 1996, Buckow, Germany; U. Müller, talks given at the Ahrenshoop Symposium, and DESY Workshop, September 25 – 28, 1996, Hamburg, Germany.
- [10] U. Müller and G.Weigt, in preparation.
- [11] T.H. Buscher, *Phys.Lett.* **B201** (1988) 466.
- [12] V.A. Fateev and A.B. Zamolodchikov, *Zh.Eksp.Teor.Fiz.* **89** (1985)380.